

Fig. 1




Fig. 2

$$C=\frac{P^2}{A} \quad or \quad C=\frac{P^2}{4\pi A} \quad or \quad C=1-\frac{4\pi A}{P^2} \quad or \quad C=\frac{A_1}{A}$$

Fig. 3

$$P=N_e+\sqrt{2}\,N_o$$

Fig. 4

$$FF=\frac{\left[\sum\limits_{n=-N/2+1}^{N/2}\|NFD(n)\|/|n|\right]}{\left[\sum\limits_{n=-N/2+1}^{N/2}\|NFD(n)\|\right]}$$

Fig. 5

$$NFD(n)=\left\{\begin{array}{ll}0;&n=0\\S(n)/S(1);&n=1,2,...,N/2\\S(n+N)/S(1);&n=-1,-2,...,-N/2+1\end{array}\right.$$

Fig. 6

$$S(n)=\frac{1}{N}\sum_{k=0}^{N-1}s(k)\exp[-j2\pi nk/N],\,n=0,1,2,\ldots,N-1,$$

Fig. 7

$$m_p=\frac{1}{N}\sum_{i=1}^N[z(i)]^p$$

Fig. 8

$$\mu_p=\frac{1}{N}\sum_{i=1}^N[z(i)-m_1]^p$$

Fig. 9

$$M=\frac{(\mu_4)^{1/4}}{m_1}-\frac{(\mu_2)^{1/4}}{m_1}$$

Fig. 10

$$\varepsilon=\frac{(m_{2,0}-m_{0,2})^2+4m_{1,1}^2}{(m_{2,0}+m_{0,2})^2}$$

Fig. 11

$$m_{pq} = \sum_x \sum_y x^p y^q f(x, y)$$


Fig. 12

$$S = \mu_{0,2} + \mu_{2,0}$$

Fig. 13


$$\mu_{pq} = \sum_x \sum_y (x - \bar{x})^p (y - \bar{y})^q f(x, y)$$

Fig. 14



BEST AVAILABLE COPY


Fig. 15 (a)



BEST AVAILABLE COPY

Fig. 15(b)

BEST AVAILABLE COPY



BEST AVAILABLE COPY

Fig. 15(c)




Fig. 16




Fig. 17





Fig. 18



BEST AVAILABLE COPY

Fig. 19




Fig. 20




Fig. 21




Fig. 22




Fig. 23




Fig. 24




Fig. 25




Fig. 26

